## 22 The Existence of Parallel Lines

The concept of parallel lines has led to both the most fruitful and the most frustrating developments in plane geometry. Euclid (c. 330-275 B.C.E.) defined two segments to be parallel if no matter how far they are extended in both directions, they never meet.

The history of the parallel postulate is fascinating. In fact, many mathematicians attempted to prove the Fifth Postulate, and some thought they had succeeded.

We shall see in this chapter that the axioms which we have adopted so far (and which are a refinement of those of Euclid) are sufficient only for proving the existence of parallel lines, but not the uniqueness.

**<u>Definition</u>**. (transversal). Given three distinct lines  $\ell$ ,  $\ell_1$ , and  $\ell_2$ , we say that  $\ell$  is a transversal of  $\ell_1$  and  $\ell_2$  if  $\ell$  intersects both  $\ell_1$  and  $\ell_2$ , but in different points.

**Definition**. (alternate interior angles, corresponding angles). Assume that the line  $\overleftrightarrow{GH}$  is transversal to  $\overleftrightarrow{AC}$  and  $\overleftrightarrow{DF}$  in a metric geometry and that  $\overleftrightarrow{AC} \cap \overleftrightarrow{GH} = \{B\}$  and  $\overleftrightarrow{DF} \cap \overleftrightarrow{GH} = \{E\}$ . If the points A, B, C, D, E, F, G and H are situated in such a way that

(i) A-B-C, D-E-F, and G-B-E-H, and (ii) A and D are on the same side of  $\overleftrightarrow{GH}$ 

then  $\angle ABE$  and  $\angle FEB$  are a pair of alternate interior angles and  $\angle ABG$  and  $\angle DEB$  are a pair of corresponding angles.

**<u>Theorem</u>**. Let  $\ell_1$  and  $\ell_2$  be two lines in a neutral geometry. If there is a transversal  $\ell$  of  $\ell_1$  and  $\ell_2$  with a pair of alternate interior angles congruent then there is a line  $\ell'$  which is perpendicular to both  $\ell_1$  and  $\ell_2$ .

**1.** Prove the above Theorem. [Th 7.1.1, p171]

<u>**Theorem</u></u>. In a neutral geometry, if \ell\_1 and \ell\_2 have a common perpendicular, then \ell\_1 is parallel to \ell\_2. In particular, if there is a transversal to \ell\_1 and \ell\_2 with alternate interior angles congruent, then \ell\_1 || \ell\_2.</u>** 

**2.** Prove the above Theorem. [Th 7.1.2, p172]

By above theorem, if  $\ell_1$  and  $\ell_2$  have a common perpendicular then  $\ell_1 || \ell_2$ . Is the converse true: If  $\ell_1 || \ell_2$ , do  $\ell_1$  and  $\ell_2$  have a common perpendicular? **3.** In the Poincaré Plane let  $\ell = {}_{0}L$  and  $\ell' = {}_{1}L_{1}$ . Show that  $\ell || \ell'$  but that there is no line perpendicular to both  $\ell$  and  $\ell'$ . [Ex 7.1.3, p173]


**<u>Theorem</u>**. In a neutral geometry, let  $\ell$  be a line and  $P \notin \ell$ . Then there is a line  $\ell'$  through P which is parallel to  $\ell$ .

**4.** Prove the above Theorem. [Th 7.1.4, p173]

**5.** Show that in the Poincaré Plane there is more than one line through P(3, 4) which is parallel to  $_{-5}L$ . [Ex 7.1.5, p174]

**Definition**. (Euclid's Fifth Postulate (EFP)) A protractor geometr satisfies Euclid's Fifth Postulate (EFP) if whenever  $\overrightarrow{BC}$  is a transversal of  $\overrightarrow{DC}$  and  $\overrightarrow{AB}$  with

(i) A and D on the same side of  $\overrightarrow{BC}$ (ii)  $m(\measuredangle ABC) + m(\measuredangle BCD) < 180$ 



then  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$  intersect at a point E on the same side of  $\overrightarrow{BC}$  as A and D.

**<u>Theorem</u>**. If  $\ell$  is a line and  $P \notin \ell$  in a neutral geometry which satisfies EFP, then there exists a unique line  $\ell'$  through P which is parallel to  $\ell$ .

**6.** Prove the above Theorem. [Th 7.1.6, p175]

<u>Definition</u>. (Euclidean Parallel Property (EPP)) An incidence geometry satisfies the Euclidean Parallel Property (EPP) if for every line  $\ell$  and every point P, there is a unique line through P which is parallel to  $\ell$ .

Note that EPP is a property of an incidence geometry so that the Taxicab Plane, Euclidean Plane, and  $\mathbb{R}^2$  with the max distance all satisfy EPP because they all have the same underlying incidence geometry, and it satisfies EPP. Of course, only the second is a neutral geometry.

<u>**Theorem**</u>. If a neutral geometry satisfies EPP then it also satisfies EFP.

7. Prove the above Theorem. [Th 7.1.7, p176]

**8.** In a neutral geometry if  $\angle ABC$  is acute then the foot of the perpendicular from A to  $\overrightarrow{BC}$  is an element of  $\operatorname{int}(\overrightarrow{BC})$ .

**9.** Given two lines and a transversal in a protractor geometry, prove that a pair of alternate interior angles are congruent if and only if a pair of corresponding angles are congruent.

## 23 Saccheri Quadrilaterals

In 1733 there appeared the book *Euclid Vindicated of All Flaw* by the Jesuit priest Gerolamo Saccheri. In it the author purported to prove Euclid's Fifth Postulate as a theorem. We now recognize basic flaws in his argument at certain crucial steps. However, the book was and is important in the development of the theory of parallels because it was the first to investigate the consequences of assuming the negation of Euclid's Fifth Postulate.

Despite his failure to actually prove Euclid's Postulate as a theorem, Saccheri did contribute a substantial body of correct results. Did he know about the flaws in his proof? Certainly the erroneous proofs were unlike any of the rest of his carefully reasoned development. It has been suggested that Saccheri knew what he did was fallacious and that the "proof" was included so that the Church would approve the publication of his work.

**Definition**. (Saccheri quadrilateral, lower base, upper base, lower base angles, upper base angles) A quadrilateral  $\Box ABCD$  in a protractor geometry is a Saccheri quadrilateral if  $\measuredangle A$  and  $\measuredangle D$  are right angles and  $\overline{AB} \cong \overline{CD}$ . In this case we write  $\Box ABCD$ . The lower base of  $\Box ABCD$  is  $\overline{AD}$ , the upper base is  $\overline{BC}$ , the legs are  $\overline{AB}$  and  $\overline{CD}$ , the lower base angles are  $\measuredangle A$  and  $\measuredangle D$ , and the upper base angles are  $\measuredangle B$  and  $\measuredangle C$ .

The basic approach of Saccheri (and those who followed him) was to try to prove something which turned out not to be true: that every Saccheri quadrilateral was actually a rectangle. If that were true it would not be hard to prove that EPP holds.

<u>**Theorem**</u>. In a neutral geometry a Saccheri quadrilateral  $\square ABCD$  is a convex quadrilateral.

**<u>Definition</u>**. (congruent convex quadrilaterals) Two convex quadrilaterals in a protractor **10.** In a neutral geometry, if  $\ell$  is a transversal of  $\ell_1$  and  $\ell_2$  with a pair corresponding angles congruent, prove that  $\ell_1 || \ell_2$ .

**11.** In a neutral geometry, if  $\overrightarrow{BC}$  is a common perpendicular of  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$ , prove that if  $\ell$  is a transversal of  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$  that contains the midpoint of  $\overrightarrow{BC}$  then a pair of alternate interior angles for  $\ell$  are congruent.

geometry are congruent if the corresponding sides and angles are congruent. In this case we write  $\square ABCD \cong \square EFGH$ .

<u>**Theorem**</u>. In a neutral geometry, if  $\overline{AD} \cong \overline{PS}$  and  $\overline{AB} = \overline{PQ}$ , then  $\Box ABCD \cong \Box PQRS$ .

**Corollary** In a neutral geometry if  $\square ABCD$  is a Saccheri quadrilateral then  $\square ABCD \cong \square DCBA$  and  $\measuredangle B \cong \measuredangle C$ .

<u>Theorem</u>. (Polygon Inequality). Suppose n > 3. If  $P_1, P_2, ..., P_n$ , are points in a neutral geometry then

 $d(P_1, P_n) \le d(P_1, P_2) + d(P_2, P_3) + \dots + d(P_{n-1}, P_n).$ 

<u>**Theorem**</u>. In a neutral geometry, given  $\overline{\mathbb{S}ABCD}$ , then  $\overline{BC} > \overline{AD}$ .

<u>**Theorem</u></u>. In a neutral geometry, given \exists ABCD, then \measuredangle ABD \le \measuredangle BDC.</u>** 

<u>**Theorem**</u>. In a neutral geometry the sum of the measures of the acute angles of a right triangle is less than or equal to 90.

<u>Theorem</u>. (Saccheri's Theorem). In a neutral geometry, the sum of the measures of the angles of a triangle is less than or equal to 180.

It must be remembered that above theorem is the best possible result. We have already seen an example of a triangle in  $\mathcal{H}$  in which the sum of the measures of the angles is actually strictly less than 180. In your high school geometry course you learned that the sum of the measures of the angles of a triangle was exactly 180. That result was correct because you were dealing exclusively with a geometry which satisfied EPP.

...for the rest of results and lots of interesting problems, see the lectures and the given book, pages 178-187...